

Frugal Chemoprophylaxis against COVID-19: Possible preventive benefits for the populace

John Oluwafemi Teibo ^{a,1}, Folashade Gloria Olorunfemi ^b, John Ajayi ^c, Abiodun Mohammed Adewolu ^b

^a Dept. of Biochemistry, Faculty of Basic Medical Sciences, College of Medicine, University of Ibadan, Nigeria

^b Dept. of Biochemistry, Faculty of Basic Medical Sciences, College of Medicine, University of Lagos, Nigeria

^c Dept. of Biochemistry, Federal University of Technology Nigeria

Received 22 April 2020; revised 22 March 2021; accepted 20 May 2021

Abstract

Introduction: Since the outbreak of Covid-19 around December 2019 which is rapidly expanding with confirmed cases in over 210 countries. Global incidence is on a ravaging increase and with new cases reported daily which has imposed threat to global health and economy. Some of the notable symptoms include dry cough, cold, fever and respiratory distress alongside inflammation.

Methods: In order to curtail the community, spread and reduce the incidence of new cases of Covid-19 we reviewed some medicinal foods, leaves and spices that have chemoprophylaxis and potent pharmacological activities.

Results and Discussion: The anti-inflammatory, anti-viral, anti-bacterial, antioxidant, anti-pyretic and inhibition of viral replication properties against the SARS-COV-2 (Severe acute respiratory syndrome coronavirus 2) symptoms and elucidate their bioactive compounds with their mechanism of action and suggest possible preventive roles they play in the abating the spread of the Covid-19 and reduce the cases globally in a bid of raising the immune system function and possible abate the development of symptoms and general health function.

Conclusion: In this present review, we presented the preventive potentials of some medicinal foods, natural products, and their role in abating the community spread of Covid-19 globally as well as the enhancement of immunity.

Keywords: Frugal Chemoprophylaxis; COVID-19; Medicinal foods; Bioactive compounds; Pharmacological properties; Mechanism of action; Preventive benefits.

1. Introduction

The outbreak of a novel coronavirus disease; a respiratory syndrome, referred to as COVID-19, was first recognized in Wuhan, China, in December 2019. The causative agent for this deadly viral infection is a coronavirus known as SARS-CoV-2. COVID-19 is demonstrated by fever, dry cough, persistent pressure in the chest, and shortness of breath (Huang et al., 2020; CCDC, 2020). There are several symptoms of coronavirus infection, such as sore throat, running nose, cough, sneezing, fever, viral conjunctivitis, loss of smell and taste and severe pneumonia (Chen et al., 2020; Li et al., 2020; Xia et al., 2020) Sneezing, runny nose, and symptoms similar to the common cold are observed in only 5% of patients. The novel coronavirus-related pneumonia COVID-19 has continued to disseminate, with the current case count close to 75,704,857 cases, and more than 1,690,061 deaths according to the World Health Organization (WHO) as of 21st December, 2020 (WHO, 2020).

The mortality rate of COVID-19 is about 7% which varies by countries and regions is less than that of SARS (severe acute respiratory syndrome), which has a mortality rate of 9.6%, and less than that of MERS (Middle East respiratory syndrome), up to 34.4% deaths. Individuals already suffering from

¹Corresponding author. Email addresses: teiboluwafemi@gmail.com Phone: +2348133577788 (J.O. Teibo)

cardiovascular disease, hypertension, respiratory disease, or diabetes are at a high risk of mortality. Age and gender-specific variations in the death rate have also been reported (WHO, 2020). The disease became pandemic within a few months of its emergence, indicating a high transmission ability as compared with SARS and MERS (Munster et al., 2020; Guan et al., 2020).

COVID-19 can last for up to 6 weeks depending upon the individual's immunity and the disease intensity. A variable incubation period has been reported for the infection to establish completely; a second exposure to the viral inoculum may decrease the incubation time (Munster et al., 2020). An incubation time of 3 to 27 days (average 14 days) has been reported by different sources (WHO, 2020; Bai et al., 2020; Lauer et al., 2019). This incubation period is considerably longer than that required by SARS or MERS (Lessler et al., 2009; Backer et al., 2020). SARS-CoV-2 has an airborne route of transmission, whereby small aerosols spread in the surrounding air by the coughing and sneezing of infected individuals. These fine airborne droplets, harboring viral particles, can be directly inhaled by nearby healthy individuals (Liu et al., 2020). The viral particles can stick to the fingertips and invade the healthy individuals by contact of contaminated hands with the nose, eyes, or mouth. Hence, hand hygiene is an expedient precaution to reduce SARS-CoV-2 transmission (Yang, 2020). The infection mechanism of both the SARS and COVID-19 viruses involve an interaction with the angiotensin-converting enzyme 2 (ACE2) and cleavage of the viral spike protein by a serine protease (Zhao et al., 2019; Prabakaran et al., 2004).

To have a better understanding of respiratory infectious disease transmission for pathogenesis and epidemiological spread of disease, a model for respiratory emissions was established and it was found that droplets containing the virus can be as small as 1 micron and a multiphase turbulent gas cloud from a human sneeze exhibited the property to travel great distance (7–8 m) (Bourouiba, 2020). This suggests that the gas cloud with its pathogen payload can span a certain space in a few seconds. The pathophysiology of SARS-CoV-2 infection closely resembles that of SARS-CoV infection, with aggressive inflammatory responses strongly implicated in the resulting damage to the airways (Wong et al., 2004). Therefore, disease severity in patients is due to not only the viral infection but also the host response.

Giving a high rate of community spread, there is a need to change the public health policy from containment to mitigation of transmission amongst community, and determine the extent to which mild disease is contagious in the community, particularly among less vulnerable young adults for acquisition of SARS-CoV-2 infection (Spellberg et al., 2020). The constant and rapid spread of novel coronavirus SARS-CoV-2 and its ability to disseminate from human to human has prompted researchers to identify existing medicinal foods that are frugal and could have suitable preventive benefits on the novel coronavirus-related pneumonia COVID-19.

Chemoprophylaxis refers to the administration of medication or natural substance for the purpose of preventing a disease or infection. The choice of chemoprophylaxis agent is dependent on if the benefits outweighs the potential harm and if it is cost effective.

Since these medicinal foods possess bioactive compounds with potent pharmacological activities such as anti-inflammatory, anti-viral, anti-bacterial, antioxidant, anti-pyretic, and inhibition of viral replication properties against the SARS-COV-2 symptoms, therefore in this review, we elucidated their various bioactive compounds with their mechanism of actions. Again, we suggest possible preventive roles they play in abating the spread of the Covid-19 and reducing the cases globally perhaps by bolstering the immune system.

2. Medicinal Foods, Leafs and Spices

These are foods that do not only provide nutrient for the body but also confers medicinal and health promoting benefits. There are many examples of these medicinal foods. In this review, we would be examining foods, leafs and spices like onion, bitter cola, bitter leaf, alligator pepper, turmeric, ginger, garlic and nutrients like zinc, ascorbic acid.

2.1 Onion (*Allium Liliaceae*)

Onions are members of the *Allium* genus of flowering plants that also includes garlic, shallots, leeks and chives. These vegetables contain various vitamins, minerals and potent plant compounds that have been shown to promote health in many ways. In fact, the medicinal properties of onions have

been recognized since ancient times, when they were used to treat ailments like headaches, heart disease and mouth sores (Nemeth and Piskula, 2007).

2.1.1 Biochemical composition of Onion

Onions (Figure 1) contain 89% water, 1.5% protein, and vitamins B₁, B₂, and C, along with potassium and selenium. Polysaccharides such as fructosans, saccharose, and others are also present, as are peptides, flavonoids (mostly quercetin), and essential oil. Methods for the qualitative assessment of the flavonoids have been detailed, and quercetin glycosides have been shown to be heat stable and transferable to cooking water (Lanzotti, 2006).



Fig. 1: Picture of Onion adopted from the internet (www.shutterstock.com)

Onion contains numerous sulfur compounds, including thiosulfinates; cepaenes; S-oxides; S-dioxides; mono-, di-, and trisulfides; and sulfoxides. When the onion bulb is minced or crushed, cysteine sulfoxides are released from cellular compartments and make contact with the enzyme alliinase from the adjacent vacuoles. Hydrolysis results in the release of reactive intermediate sulfenic acid compounds and then the various sulfur compounds which can be utilized as a potential preventive property on the novel Covid-19 virus (Arnault and Auger, 2006).

2.1.2 Potential Mechanisms of Pharmacological Uses of Onion and its Potential Use as Chemoprophylaxis against Covid-19

Anti-inflammatory effects: Animal and in vitro data shows that compounds derived from onion have exerted anti-inflammatory and antihistamine effects in vitro and in animal models (Griffiths et al., 2002).

Antimicrobial effects: In vitro studies have shown onion possesses antibacterial (example is *Helicobacter pylori*), antiparasitic, and antifungal activity (Elnima et al., 1983). Clinical applications for this activity have not been determined, and use as a food preservative is limited by the strong odor and instability of sulfur compounds (Griffiths et al., 2002).

2.1.3 Dosing

An average daily dose of 50 g of fresh onion, or 20 g of dried onion has been suggested for a variety of uses (Nathan and Scholten, 1999; WHO, 1999).

2.1.4 Possible Adverse Reactions of Onion

Ingestion of onion and onion extract appears to be relatively safe (Nathan and Scholten, 1999; WHO, 1999). Excessive consumption of onion has been associated with GI upset (burning sensation and diarrhea), flatulence, and changes in intestinal flora (Suleira et al., 2015). Onion seeds have been reported as occupational allergens, but frequent contact with onion rarely causes allergic reaction

(Navarro et al., 1995). Certain sulfur compounds, such as propanethial-S-oxide, escape from onion in vapor form and hydrolyze to sulfuric acid when cut, causing eye irritation and lacrimation (Chan et al., 2020).

2.2 Bitter Kola (*Garcinia kola*)

Garcinia kola (bitter kola) is a dicotyledonous plant belonging to the family of plants called Guttiferae. It is a perennial crop growing in the forest, distributed throughout West and Central Africa (Iwu, 1993). *G. kola* is also found distributed in the forest zone of Sierra Leone, Ghana, Cameroon and other West African countries. In Nigeria, it is common in the South Western States and Edo State (Otor et al., 2001).

It is a medium sized evergreen tree, about 15-17m tall and with a fairly narrow crown. The leaves are simple, 6-14cm long and 2-6cm across, shiny on both surfaces and spotted with resin glands. The small flowers are covered with short, red hairs (Iwu, 1993). The fruit is a drupe of 5-10cm in diameter and weighs 30-50g. It is usually smooth and contains a yellow-red pulp. The fruit changes color during maturation from green to orange, and each fruit contains 1-4 seeds (Juliana et al., 2006).

Garcinia kola (Figure 2) has been referred to as a “wonder plant” because every part of it has been found to be of medicinal importance (Dalziel, 1937). It is also called bitter cola, male kola due to the reported aphrodisiac properties. It is commonly called “Orogbo” in Yoruba language, *Aku ilu* in Igbo language and *Namijin goro* in Hausa language (Dalziel, 1937).



Fig. 2 : Picture of bitter kola adopted from the internet (www.9jafoods.com)

2.2.1 Biochemical constituents

The phytochemical compounds isolated from *G. kola* include tannins, saponins, alkaloids, cardiac glycosides (Ebana et al., 1991). Other phytochemical compounds isolated from *G. kola* seeds are biflavonoids such as kolaflavone and 2-hydroxybi-flavonols. Two new chromanols, garcioic and garcinal, together with tocotrienol were reported isolated from *G. kola* (Terashima et al., 2020; Morabandza et al., 2013) had also determined the chemical composition of *Garcinia kola* Heckel (Clusiaceae) mesocarp.

2.2.2 Traditional Uses and Medicinal Values

Garcinia kola is chewed extensively in Southern Nigeria as a masticatory and it is readily served to visitors, especially among the Igbo tribe in Eastern Nigeria, as a sign of peace and acceptance of visitors. The root of the plant is used as favourite bitter chew-sticks in West Africa (Otor et al., 2001).

The stem bark is used in folklore remedies as a purgative among the natives of Eastern Nigeria and the latex is externally applied to fresh wounds to prevent sepsis, thereby assisting in wound healing. It is also popular among the people of Nigeria for nervous alertness and induction of insomnia. *Garcinia kola* is highly valued for medicinal use. This plant has been referred to as a wonder plant because every part of it has been found to be of medicinal importance (Dalziel, 193). The seeds are chewed as an aphrodisiac or used to cure cough, dysentery, chest colds, liver disorders, diarrhoea, laryngitis, bronchitis, and gonorrhea (Adesina et al., 1995). The seed is used to prevent and relieve colic; it can also be used to treat headache, stomach ache and gastritis (Ayensu, 1978). It has also been reported for

the treatment of jaundice, high fever, and as purgative (Iwu, 1989). In Sierra Leone, the roots and bark are taken as a tonic for sexual dysfunction in men.

The bark is also added into palm wine to improve its potency (Iwu et al., 1990). Traditional medicine practitioners in Nigeria, particularly in the Ogoni area use a decoction of *Garcinia kola* stem bark for the treatment of dysmenorrhoea, fever, inflammation and burns (Adesina et al., 1995). "Bitter kola is anti-poison and helps to detoxify the system, it has the ability to repel evil men and spirits, it could sound superstitious but it works (Iwu, 1989).

2.2.3 Possible Mechanism of action of Garcinia kola as Chemoprophylaxis agent against Novel Covid-19

2.2.3.1 Anti-microbial properties

Adegboye, et al. (Adegboye et al., 2008) had investigated the in vitro antimicrobial activities of crude extract of *Garcinia kola* against some bacterial isolates comprising of both Gram-positive and Gram negative organisms. In another study, the antimicrobial interaction between *Garcinia kola* seed (GKS) and gatifloxacin (GAT), a fourth generation fluoroquinolone was evaluated by a modification of the checkerboard technique using *Bacillus subtilis* and *Staphylococcus aureus* as the test organisms (Ofokansi et al., 2008). The results showed that the effect of combination of the methanol extract of GKS with gatifloxacin was dependent not only on the ratio of combination but also on the test organism employed for the evaluation. Overall, the combined antimicrobial effect of the interaction between GKS and gatifloxacin was predominantly synergistic against *B. subtilis* (Ofokansi et al., 2008). The antimicrobial activity of five solvent extracts of *Garcinia kola* seeds had also been investigated against 30 clinical strains of *H. pylori* and a standard control strain, NCTC 11638, using standard microbiological techniques (Collise et al., 2011). In vitro anti-*Vibrio* activities of methanol and aqueous extracts of *Garcinia kola* seeds against 50 *Vibrio* isolates obtained from wastewater final effluents in the Eastern Cape Province, South Africa were also investigated (Penduka et al., 2011). The bioactivity of *G. kola* seeds was also assessed on *Streptococcus pyogenes*, *Staphylococcus aureus*, *Plesiomonas shigelloides* and *Salmonella typhimurium* (Seanego et al., 2012). *Trichomona* cidal effects of *G. kola* nuts were also investigated (Gabriel and Emmanuel, 2011). Esimone et al. (Esimone et al., 2002) also investigated the effect of *Garcinia kola* seed extract (100 mg/kg) on the pharmacokinetic and antibacterial effects of ciprofloxacin hydrochloride (40 mg/kg).

2.2.3.2 Hepatoprotective and Anti-oxidant activities

The hepatoprotective effect of *Garcinia kola* seed extract against paracetamol induced hepatotoxicity had been investigated in rats which their results showed that the hepatoprotective effect of the extract may be due to inhibition of cytochrome P-450 which normally converts paracetamol to the toxic intermediate N-acetyl-p-benzoquinoneimine (NAPQI) (Akintonwa and Essien, 1990). The protective effects of *Garcinia kola* against a dose of Carbon-Tetrachloride (CCl₄)-induced liver damage in experimental rats were also investigated which from the results observed there was a significant decrease in liver marker enzymes and lipid peroxides (Matthew and Blessing, 2007). Antioxidant potentials of five fractions (ME1–ME5) of methanolic extract of *Garcinia kola* seeds was also studied in which moreoflavone and GB2a very abundant compounds in the seeds of this plant, could be responsible, at least in part, for the antioxidant potential of its crude extract (Tebekeme, 2009).

2.2.4 Other studies on Garcinia kola and its health benefits

Anti-ulcer potential and proton pump inhibitory activity of kolaviron (KV) isolated from *Garcinia kola*. Heckel had been evaluated using different ulcer models and was suggested to emerge as a potent anti-ulcer compound (Onasanwo et al., 1990). *Garcinia kola* (Heckel) seed extract evaluated in albino Wistar rats possess significant anti-pyretic activity, which justified its ethnomedicinal use (Kakjing et al., 2014). *Garcinia kola* 0.5% aqueous solution eye drop significantly reduces intraocular pressure (IOP) as compared to baseline (Adebukunla et al., 2010). Comparative study on the efficacy of *Garcinia kola* in reducing some heavy metal accumulation in liver of Wistar rats was also carried out. *Garcinia kola* has the highest hepatoprotective effect to Cd followed by Hg and least protection against

Pb toxicity in rats and its administration was beneficial in reducing heavy metal accumulation in the liver (Nwokochaa et al., 2011).

Administration of *Garcinia kola* for a period of six weeks in rabbits elicited no observable histopathological effects on the histology of the liver (Charity et al., 2012). *Garcinia kola* had been shown to enhance erythropoiesis in rabbits and rats and as well has no long term significant toxicological implication (Unigwe and Nwakpu, 2009). In another study, *Garcinia kola* significantly reduced tissue damage induced by lipopolysaccharide (LPS) (Okoko and Ndoni, 2009). Daily administration of *Garcinia kola* (*G. kola*) in growing Wistar rats for a period of 70 days showed a depressive effect on appetite and water intake with resultant poor feed utilization efficiency and mass gain of rats in a dose dependent manner. Plasma alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities were elevated ($P < 0.05$) but histological examinations of liver, heart and lungs of experimental rats revealed no alterations. The extract enhanced sexual interest (libido) of the male rats but did not necessarily improve their fertility rate (King et al., 2006).

2.3 Zinc

Zinc essentiality was established in 1869 for plants, in 1934 for experimental animals and in 1961 for humans (King et al., 2006). A syndrome of anemia, hypogonadism and dwarfism was reported in a 21-year-old Iranian farmer in 1961 who was subsisting on a diet of unrefined flat bread, potatoes, and milk (Prasad et al., 1963). Shortly after, a similar syndrome was observed in Egyptian adolescents who had similar dietary history to that of the Iranians, mainly subsisting on bread and beans (Sandstead et al., 1967).

2.3.1 Biochemical and Physiologic Functions: Although, zinc-dependent biochemical mechanisms in physiologic functions have received extensive study, clear relationships have not been fully established. Zinc is ubiquitous within cells in contrast to iron, which is contained in defined cellular components and has defined physiological roles. The role of zinc in biology can be grouped into three general functional classes, namely catalytic, structural and regulatory functions (Cousins et al., 1996).

Zinc is released from food as free ions during digestion. These liberated ions may then bind to endogenously secreted ligands before their transport into the enterocytes in the duodenum and jejunum (Turnlund et al., 1984). Specific transport proteins may facilitate the passage of zinc across the cell membrane into the portal circulation. With high intakes, zinc is also absorbed through a passive paracellular route. The portal system carries absorbed zinc directly to the liver, and then released into systemic circulation for delivery to other tissues. About 70% of the zinc in circulation is bound to albumin, and any condition that alters serum albumin concentration can have a secondary effect on serum zinc levels. Although, serum zinc represents only 0.1% of the whole body zinc, the circulating zinc turns over rapidly to meet tissue needs (Tubek, 2007).

2.3.2 Possible Mechanism of action of Zinc as Chemoprophylaxis agent against Novel Covid-19

Diarrhea: Diarrhea is characteristically, although not inevitably, a prominent feature of acrodermatitis enteropathica. (Hambidge, 2001). Plausible explanations for a link between zinc deficiency and diarrhea include impairment of the immune system and of intestinal mucosal cell transport (Ghishan, 1984). A causal relationship between zinc deficiency and diarrhea is indicated by the beneficial effects of zinc supplements and concurrent increase in growth velocity (Brown et al., 1998).

Pneumonia: Community zinc supplementation studies in children have demonstrated a substantial and statistically significant reduction in the prevalence of pneumonia in developing countries (Bhutta et al., 1999), this makes its effective in treatment of pneumonia complications which results from Covid-19.

Malaria: It is uncertain to what extent oral supplementation with zinc can reduce episodes of malaria in endemic areas. According to some studies, malaria also appears to be reduced by zinc supplementation (Shankar et al., 2000) However, there are studies showing no effect of zinc supplementation against malaria (Veenemans et al., 2011). Further studies are required to establish this effect.

2.4 Alligator Pepper (*Aframomum melegueta*)

This plant is a member of the ginger family (Zingiberaceae) and is widely cultivated in tropical areas of Africa, mostly in West Africa. *Aframomum melegueta* (Grains of Paradise) has names such as grains of paradise, Atare (in Yoruba), chitta (Hausa), is one seed with many healing power and of great benefits to mankind (Figure 3).

The seeds of the plant are used to flavor foods and as components of traditional African medicine. Alligator pepper has been reported to have anti-ulcer, anti-microbial, anti-nociceptive, anti-plasmodial, hepatoprotective and anticancer activities (El-Halawany et al., 2014).



Fig. 3: *Aframomum melegueta* (Alligator pepper) adopted from internet (www.e-time.com)

Traditionally, *Aframomum Melegueta* is mixed with other herbs for the treatment of common ailments such as body pains, diarrhoea, sore throat, catarrh, congestion and rheumatism in West Africa, Nigeria (Ajaiyeoba and Ekundayo, 1999). Diabetes, a major underlying disease of people afflicted with covid-19 might also be combated by *Aframomum melegueta*. A myriad of compounds found in *Aframomum melegueta* such as 6-paradol, 6-shagaol, 6-gingerol, oleanolic acid and acarbose exert an anti-diabetic effect by inhibiting enzymes such as α -amylase and α -glucosidase. These enzymes are responsible for digestion and break down of carbohydrates and polysaccharides from food into simple sugars to increase blood glucose levels. Among the compounds, 6-gingerol and oleanolic acid are more effective in inhibiting the enzymes (Mohammed et al., 2017). *Aframomum melegueta* is suitable for consumption by diabetic patients; its consumption will help a diabetic patient stay healthy (Venugopal and Liu, 2012).

The ethanolic seed extract and stem bark of *Aframomum melegueta* contains phytochemicals such as tannins, saponins, flavonoids, steroids, terpenoids, cardiac glycosides and alkaloids that possess antimicrobial and anti-inflammatory effects (Doherty et al., 2010; Okwe, 2005; Okoli et al., 2007). These compounds which are natural antioxidant are believed to scavenge free radicals and offer protections against viruses, allergens, microbes, platelet aggregation, tumors, ulcers and hepatotoxins (chemical liver damage) in the body. To support this claim of anti-inflammatory properties of *Aframomum melegueta*, Ilic et al. (Ilic et al., 2014) reported that ethanolic *Aframomum melegueta* extract inhibited cyclooxygenase-2 (COX-2). Compounds that inhibit COX-2 activity are capable of reducing inflammatory responses. The most active COX-2 inhibitory compound in the *Aframomum melegueta* extract was [6]-paradol, while [6]-shogaol was found to inhibit expression of a pro-inflammatory gene, interleukin-1 beta (IL-1 β) (Osuntokun et al., 2017).

2.5 Turmeric (*Curcuma longa*)

Turmeric (*Curcuma longa* L.) belongs to the family of ginger (Zingiberaceae) and natively grows in India and Southeast Asia. The plant's rhizomes contain several secondary metabolites including curcuminoids, sesquiterpenes, and steroids (Omola et al., 2017), with the curcuminoid curcumin being the principal component of the yellow pigment and the major bioactive substance. It has been used as traditional medicine from ancient time, especially in Asian countries. Plants as a rich source of phytochemicals with different biological activities including antiviral activities are in interest of scientists (Jassim and Naji, 2003). It has been demonstrated that curcumin as a plant derivative has a wide range of antiviral activity against different viruses. Inosine monophosphate dehydrogenase (IMPDH) enzyme due to rate-limiting activity in the de novo synthesis of guanine nucleotides is suggested as a therapeutic target for antiviral and anticancer compounds. Among the 15 different polyphenols, curcumin through inhibitory activity against IMPDH effect in either noncompetitive or competitive manner is suggested as a potent antiviral compound via this process (Dairaku et al., 2010).

Furthermore, protein molecules encoded by the SARS-CoV-2 genome are potential targets for chemotherapeutic inhibition of viral infection and its replication. These intriguing targets include the spike protein (S), which mediates the entry of the virus, the SARS-CoV-2 main protease (3CL protease), the NTPase/helicase, the RNA-dependent RNA polymerase, the membrane protein (M) required for virus budding; the envelope protein (E) which plays a role in coronavirus assembly (Stadler et al., 2003; Lai, 2005; Holmes, 2003) and the nucleocapsid phosphoprotein (N) that relates to viral RNA inside the virion and possibly other viral protein-mediated processes (Gallagher and Buchmeier, 2001). With such a drastic increase in molecular and biochemical information about various components of the SARS-CoV-2 and their cellular targets, it is important and timely to again evaluate anti-SARSCoV-2 activities of various turmeric phytochemicals. It is to be noted that in vitro studies have positively demonstrated effective antiviral properties of curcumin against enveloped viruses such as Dengue virus (DENV), influenza virus and emerging arboviruses like the Zika virus (ZIKV) or chikungunya virus (CHIKV) (Chen et al., 2013; Munaya 2013) which are similar to the SARS-CoV-2 an enveloped virus.

2.6 Bitter Leaf (*Vernonia amygdalina*)

Vernonia amygdalina (Del.) commonly called bitter leaf is the most widely cultivated species of the genus *Vernonia* which has about 1,000 species of shrubs (Munaya 2013). Bitter leaf is a seedless plant, green in colouration with a characteristic odour and bitter taste. It has different names such as 'Omjunso' in East Africa especially Tanzania, 'Ewuro' in Yoruba, Etidot (Ibibio), Ityuna (Tiv), Oriwo (Edo), Chusa-doki Shiwaka (Hausa), and 'Omubirizi' in south-western Uganda.

Vernonia amygdalina is known as food and medicinal plants used in Asia and Africa (West Africa) due to its pharmacological effects which include antioxidant, anti-inflammatory, anti-diabetes, anticancer, anti-malaria etc. One major properties of considering *Vernonia amygdalina* in the management of covid-19 is its antioxidants property. Antioxidants inhibit deleterious effects of free radicals that are capable of deteriorating lipid biomembranes in the human body. Researchers in the field of medical sciences have observed free radical scavenging ability and antioxidant property in *Vernonia amygdalina*.

Vernonia amygdalina has been documented to possess antioxidant properties which correlates to its medicinal properties. Antioxidant activities of bioactive compounds isolated from *Vernonia amygdalina* leaves have been established from various studies (Igile et al., 1994; Farombi and Owoeye, 2011; Ho et al., 2012; Udochukwu et al., 2015). Farombi and Owoeye (Farombi and Owoeye, 2011) observes phytochemicals compounds such as saponins and alkaloids, terpenes, steroids, coumarins, flavonoids, phenolic acids, lignans, xanthones, anthraquinones, edotides, sesquiterpenes extracted and isolated from *Vernonia amygdalina* to elicit various biological effects in humans including cancer chemoprevention. The chemoprophylaxis properties of *Vernonia amygdalina* was attributed to their abilities to scavenge free radicals, induce detoxification, inhibit stress response proteins and interfere with DNA binding activities of some transcription factors.

2.7 Black Seeds (*Nigella sativa*)

Nigella sativa, commonly known as black seed is native to Southern Europe, North Africa and Southwest Asia and it is cultivated in many countries in the world like Middle Eastern Mediterranean region, South Europe, India, Pakistan, Syria, Turkey, Saudi Arabia (Khare, 2004).

N. sativa is regarded as a valuable remedy for various ailments, the seeds, oil and extracts have played an important role over the years in ancient Islamic system of herbal medicine. Bukhari reported that Holy Prophet Muhammad (peace be upon him) told “There exists, in the black grains, health care of all the diseases, except death” (Gheznavi et al., 1988).

Many studies have been carried out to affirm the acclaimed medicinal properties emphasized on different pharmacological effects of *N. sativa* seeds such as antioxidant, anti-bacterial, anti-tussive, anti-lipid peroxidative (Hosseinzadeh et al., 2007; Hosseinzadeh et al., 2008), gastroprotective, anti-anxiety (El-Abhar et al., 2003; Bin et al., 2014), anti-inflammatory, anti-cancer, immunomodulatory, anti-tumor properties (Shafi et al., 2009; Salem, 2005; Majdalawieh et al., 2010) and hepatoprotective effect (Khan, 1999).



Fig. 4: *Nigella sativa* plant and seeds adopted from Sabana et al. (2012)

It should be noted that the seeds of *N. sativa* are the source of the active ingredients of this plant. Thymoquinone (TQ), dithymoquinone (DTQ), which is believed to be nigellone, thymohydroquinone (THQ), and thymol (THY), are considered the main phytochemicals of the seed (Omar et al., 1999). *N. sativa* seeds (Figure 4) contain other ingredients, including nutritional components such as carbohydrates, fats, vitamins, mineral elements, and proteins, including eight of the nine essential amino acids (Chun et al., 2002; Correa et al., 1986). The potential immunomodulatory and immunotherapeutic potentials of *N. sativa* seed active ingredients, most especially TQ will show a promising therapy in managing covid-19 virus.

1.8 Ginger (*Zingiber Officinale Roscoe*)

Ginger, with the botanical name *Zingiber Officinale Roscoe* a commonly used spice and belongs to the family Zingiberaceae (Singh and Singh, 2019). Its local names are Ata-ile, Jinja, and Cithar among the Yoruba, Igbo and Hausa folks of Nigeria respectively (Nwauzoma and Dappa, 2013). Commonly, ginger can be found in subtropical and tropical Asia, Africa, Far East Asia, China, and India (Tanaka et al., 2015). It can be used as a component in curry powder, sauces, ginger bread, and ginger-flavoured carbonated drinks and also in the preparation of dietaries for its aroma and flavour (Durak et al., 2015). Investigations, reveal that ginger possesses several biological properties such as anti-inflammatory and anti-oxidative, antimicrobial, neuroprotective, cardiovascular protective, antidiabetic and anti-nausea, antiemetic activities and chemo-protective effects (Chung, 2019; Beristain-Bauza et al., 2019; Sahin et al., 2019; Ilkhanizadeh et al., 2016; Mele, 2019; Amar et al.,

2019). Its main bioactive components are 6-gingerol, 6-shogaol which are phenolics, thus accounting for its various bioactivities (Sahardi and Makpol, 2019).

Till date, there are no antiviral therapeutics that specifically target human corona viruses, and thus completely eradicating the pandemic. Even though several potential vaccines have been developed, such as recombinant attenuated viruses, live virus vectors, or individual viral proteins expressed from DNA plasmids (Fehr and Pelman, 2015). However, none have been yet approved for use. The limitation with the use of vaccines is a propensity of the viruses to recombine, thereby posing a problem of rendering the vaccine useless and potentially increasing the evolution and diversity of the virus into a more virulent form (Wang et al., 1993). It therefore implies that a preventive approach could be a better option.

2.8.1 Possible Mechanism of action of Ginger against Corona virus

Virus–receptor interactions play a key regulatory role in viral host range, tissue tropism, and viral pathogenesis (Maginnis, 2018). Many α -coronaviruses utilize amino-peptidase N (APN) as their receptor (Reguera et al., 2012), SARS-CoV and HCoV-NL63 use angiotensin- converting enzyme 2 (ACE2) as their receptor (Li et al., 2020), MHV enters through CEACAM1 (Li, 2014), and the recently identified MERS-CoV binds to dipeptidyl-peptidase 4 (DPP4) to gain entry into human cells (Raj et al., 2013). Following receptor binding, the viruses then proceeds into the cytosol of the host. A possible mechanism of preventing access of the viruses into the host's cell could be the inhibition of microbial binding to cellular receptors. In the antimicrobial mechanism of ginger, inhibition of receptor binding has been identified as a major mechanism.

2.8.2 Angiotensin converting enzyme-2 (ACE-2) binding

Hitherto, ACE-2 has been associated with an obscure function, not until in recent times with the widespread of the corona virus disease (COVID-19). It is now established as a protein being recognized by various corona viruses to gain entry into hosts cell (Li et al., 2020) suggesting it to be a therapeutic target in the treatment of the disease. We propose that both ACE receptor antagonism and agonism can be an effective mechanism of ginger in its chemopreventive strategy against corona viruses. These claims are supported by a recent report of Cava et al. (Cava et al., 2020), who demonstrated through computational methods that anti-ACE 2 compounds could be effective in blocking the replication of SARS-CoV variant. Moreover, Huentelman et al. (Huentelman et al., 2004) also used docking methods to predict the inhibitory ability of small molecules against ACE 2 enzymes, and possibly inhibiting SARS corona virus S protein mediated cell fusion, and high binding affinity was scored. In addition, different bioactive compounds in ginger have been shown to have different effects on ACE2. One mechanism is through ACE 2 inhibition. A study by Alu'datt et al. (Alu'datt et al., 2016) in the bid to evaluate the anti-hypertensive activity of ginger, demonstrated that its methanolic extract has excellent inhibitory activity on ACE 2 at an extraction temperature and time of 40°C and 6 hours respectively. To further corroborate our hypothesis, 3-3'-digallate, a phenolic compound present in ginger was found to interact with ACE2 receptor in a recent computational modelling studies (Zhang et al., 2019). Taken together, we suggest that the intake of ginger, might be beneficial for both prevention and treatment of the pandemic COVID-19 corona virus. Below (Figure 5) is a proposed inhibitory mechanism of action of Ginger against ACE-2 in its chemoprophylaxis strategy.




Fig. 5: Ginger is able to inhibit ACE-2, thereby preventing the binding of the corona virus to the protein.

1.9 Garlic (*Allium sativum* L.)

Garlic (*Allium sativum* L.) is a widely consumed spice in the world, and it contains diverse phytochemicals such as organosulfur compounds, saponins, phenolic compounds, and polysaccharides (Diretto et al., 2017; Szychowski et al., 2018; Bradley et al., 2016; Wang et al., 2018). However, its major bioactive compounds (figure 6) are the organosulfur compounds including allicin, alliin, diallyl sulfide, diallyl disulfide, diallyl trisulfide, ajoene, and S-allyl-cysteine (Shang et al., 2019). It has been reported that garlic and its constituents exhibit different bioactivities such as anti-carcinogenic, antioxidant, anti-diabetic, renoprotective, anti-atherosclerotic, antibacterial, antifungal, and antihypertensive activities.




Fig. 6: Structures of some bioactive compounds in garlic excerpted from Shang et al. (2019)

From the foregoing, worthy of relevance to our subject matter among the bioactivities of garlic is its antiviral properties. We suggest that this could be due in part to its anti-inflammatory and antioxidative abilities. The antiviral potential of garlic extracts has been demonstrated against influenza virus and entero-virus, HIV virus, herpes simplex type 2, dengue virus, Infectious Bronchitis virus, Newcastle Disease virus (Sharma, 2019; Bezerra et al., 2018; Weber et al., 1992; Hall et al., 2016; Shojai et al., 2016; Harazem et al., 2019).

2.9.1 Possible Mechanisms of action of Garlic against Corona virus

Major challenges in treating viral diseases are the development of resistance in the virus against the antiviral drugs, due to the high mutation rate of the viral RNA Polymerase (Elena and Sanjuan, 2005) and some of the antiviral drugs can also induce negative side effects in the body (Bindu and Anusha, 2011). Therefore, it is necessary to find alternative therapies in natural plants and their products, out of which garlic stands out significantly, because of its robust pharmacological bioactive components as stated above. Below is a possible mechanism of action of garlic for the prevention and treatment of the human corona virus.

2.9.1.1 Blocking of the attachment of the virus with cell membranes through protein inhibition

Following receptor binding, the corona virus then fuses with the cell membrane (Fehr and Perlman, 2015). Coronaviruses are enveloped positive-stranded RNA viruses that replicate in the cytoplasm. To deliver their nucleocapsid into the host cell, they rely on the fusion of their envelope with the host cell membrane, which is mediated by the spike (S) glycoprotein, a class I fusion protein (Sandrine et al., 2012). Inhibition of this protein by peptide inhibitors is an important mechanism of antiviral drugs. These molecules could stabilize the viral envelope protein or alter its conformation, thereby blocking oligomerization or the conformational changes required for fusion (Teissier et al., 2011). Using the S-HR1 and S-HR2 sequence in the protein as a target, Liu et al. (Liu et al., 2009), Lu et al. (Lu et al., 2014) and Xia et al. (Xia et al., 2019) have previously designed and developed several potent fusion inhibitors against the Severe Acute Respiratory Syndrome corona virus (SARS-CoV), the Middle East Respiratory Syndrome (MERS-CoV) and the Human corona virus (H-coV) respectively. Antimicrobial compounds from garlic have been shown to react with viral envelope and inhibit the penetration and exponentiation of influenza virus in animals using hemagglutination (HA), methyl tetrazolium (MTT) cytotoxicity assay and RT-PCR (Mehrbood et al., 2013). Interestingly, allicin and quercetin, which are bioactive compounds of garlic were found to exhibit inhibitory action on the COVID-19 M^{pro} protease, thereby demonstrating similar binding affinity when compared to other potent peptide inhibitors such as Nelfinavir and lopinavir in an in silico study of protein-protein docking (Khaerunnisa et al., 2019). Furthermore, it was demonstrated by Weber et al. (Weber et al., 1992) that treatment of different bioactive components of garlic against different viral strains led to the inhibition of viral adsorption or penetration. Taken together, we suggest that through this mechanism, garlic could prevent the attachment of the corona virus to the host's cellular machinery.

2.10 Vitamin C (Ascorbic acid)

Vitamin C or ascorbic acid (AA), a water-soluble vitamin was first isolated in 1923 by Hungarian biochemist and Nobel laureate Szent-Gyorgyi and synthesized by Howarth and Hirst (Haworth et al., 1933). Humans cannot synthesize ascorbic acid as they lack an enzyme called gulonolactone oxidase. It plays an important role in bone formation, wound healing and the maintenance of healthy gums, metabolic functions including the activation of the B vitamins, folic acid, the conversion of cholesterol to bile acids and the conversion of the amino acid, tryptophan, to the neurotransmitter, serotonin. As an antioxidant that protects body from free radical damage (Chambial et al., 2013).

2.10.1 Therapeutic Roles

It is used as therapeutic agent in many diseases and disorders. Vitamin C protects the immune system, reduces the severity of allergic reactions and helps to fight off infections. Large doses of vitamin C have been widely used in the treatment and prevention of a large number of disorders like diabetes, atherosclerosis, common cold, infertility, cataracts, glaucoma, macular degeneration, stroke, heart diseases, neurodegenerative diseases, cancer etc.

2.10.2 Dietary Sources

Vitamin C is commonly found in fruits and vegetables. Fruits containing vitamin C, includes; Guava, red pepper, orange, grape fruit, strawberries, banana, Brussels sprouts, pawpaw, tomatoes, mango etc. (Schlueter et al., 2010). Milk, eggs, cheese, and nuts are also rich sources of vitamin C.

2.10.3 Deficiency

Ascorbic acid deficiency gives rise to the appearance of scurvy, which is more common in developed countries. Symptoms develop with plasma levels below 0.15 mg/dL. Scurvy is characterized by the presence of weakness, joint pain or skin lesions in form of petechias, gum bleeding, ease of developing bruises or delay in wound healing. The most characteristic skin manifestations are purpuric perifollicular hyperkeratotic papules and the presence of kinky hair (Valdes, 2006).

2.10.4 Toxicity

Vitamin C is generally safe and well tolerated, even in large doses. The IOM set the Tolerable Upper Intake Level for oral vitamin C ingestion at 2 g daily for adults based on gastrointestinal disturbances observed in some individuals at higher doses. High amounts of vitamin C intake have been associated with an increased risk of kidney stones, although the evidence is mixed and inconsistent. The current recommendation is to avoid vitamin C supplementation in those susceptible to kidney stone formation. Vitamin C consumed with iron could increase the risk of iron overload in susceptible individuals. Patients with these conditions should not avoid eating fruit and vegetables but limit their intake of iron instead. Vitamin C has been reported to cause hemolysis in individuals with glucose-6-phosphate dehydrogenase deficiency, but these reports have not been substantiated (Lykkesfeldt et al., 2014).

2.10.5 Possible Mechanisms of action of Vitamin C against Corona virus

2.10.5.1 Improve host's immune responses to combat diseases

Vitamin C (Ascorbic acid) has been reportedly found to exhibit many of its therapeutic functions, partly due to it been able to improve the body's immune responses through a number of mechanisms. We therefore opine that through this physiological response, it would prove to be effective as a chemo-preventive alternative to antiviral drugs against COVID 19 corona virus.

Vitamin C plays important roles in immune function and the modulation of host resistance to infectious agents, reducing the risk, severity, and duration of infectious diseases as supplementation of vitamin C was found to improve components of the human immune system such as antimicrobial and natural killer cell activities, lymphocyte proliferation, chemotaxis, and delayed-type hypersensitivity (Wintergerst et al., 2006). Vitamin C contributes to immune defense by supporting various cellular functions of both the innate and adaptive immune system. It supports epithelial barrier function against pathogens and promotes the oxidant scavenging activity of the skin, thereby potentially protecting against environmental oxidative stress (Lauer et al., 2013; Valacchi et al., 2016). Vitamin C accumulates in phagocytic cells, such as neutrophils, and can enhance chemotaxis, phagocytosis, generation of reactive oxygen species, and ultimately microbial killing (Carr et al., 2017). It is also needed for apoptosis and clearance of the spent neutrophils from sites of infection by macrophages, thereby decreasing necrosis/netosis and potential tissue damage. (Sharma et al., 2004). Based on this established premises, we suggest that Vitamin C consumption can serve as a prophylactic measure against the pandemic. It is therefore necessary that diet formulations with adequate vitamin C supplement should be given more attention by different regulatory agencies in the local and international communities, while the populace should consume more of foods that are rich in this vitamin as we have afore mentioned.

3. Conclusion

Coronavirus is a respiratory viral infection of global health concern because the virus is contagious and may cause life-threatening respiratory infection and severe pneumonia in humans. The

pathophysiology of SARS-COV-2 infection, is associated with aggressive inflammatory responses which uncontrolled inflammation inflicts multi-organ damage leading to organ failure, especially of the cardiac, hepatic and renal systems and most patients with this infection who progressed to renal failure eventually death.

In this review, some medicinal foods were elucidated which include ginger, garlic, onion, bitter kola, alligator pepper, black seed, turmeric, bitter leaf, zinc and vitamin C which possess potent pharmacological activities which include anti-inflammatory, anti-viral, anti-pyretic properties, inhibition of viral replication properties against the symptoms displayed by SARS-COV-2 infection.

These frugal medicinal foods offer great chemoprophylactic benefits and can play potent roles in the prevention and curtailing of the community spread of COVID-19 as it bolsters immunity and defence of the body system coupled with its cost effectiveness and availability features.

We hereby recommend that further research should be done to evaluate the protective roles of these medicinal foods on COVID-19 infection as well as promote the development of drugs and improved plant medicines as this outcome can prepare the global populace against subsequent global outbreak of viruses.

Acknowledgment(s)

We are grateful to God Almighty and the Chemoprophylaxis team for their input in this work.

References

- Adebukunola OA, Bernice OA, Adebayo KA, Olayinka OA, Elsie S, Kehinde O. (2010). Efficacy of *Garcinia kola* 0.5% Aqueous Eye Drops in Patients with Primary Open Angle Glaucoma or Ocular Hypertension. *Middle East Afr J Ophthalmol*. 17: 88–93.
- Adegboye MF, Akinpelu DA, Okoh AI. (2008). The bioactive and phytochemical properties of *Garcinia kola* (Heckel) seed extract on some pathogens. *Afr J Biotechnol*. 7: 3934-3938.
- Adesina SK, Gbile ZO, Odukoya OA. (1995) Survey of indigenous plants of West Africa with special emphasis on medicinal plants and issues associated with management. *The United Nations Programme on Natural Resources in Africa*; 2nd edition. 84–85.
- Ajaiyeoba EO, Ekundayo O. (1999) Essential oil constituents of *Aframomum melegueta* (Roscoe) K. Schum. seeds (alligator pepper) from Nigeria. *Flav Frag J* 14:109-111.
- Akintonwa A, Essien AR. (1990) Protective effects of *Garcinia kola* seed extract against paracetamol-induced hepatotoxicity in rats. *J Ethnopharmacol* 29: 207-211.
- Alu'datt MH, Rababah T, Alhamad MN, Gammoh S, Ereifej K, Johargy A, Kubow S, Almajwal AM, Rawashdeh M. (2016) Optimization of Phenolic Content, Antioxidant, and Inhibitory Activities of α -Glucosidase and Angiotensin Converting (AC) Enzymes from *Zingiber officinale* Z., *International Journal of Food Properties*, 19:6: 130-1316.
- Amar U, Omprakash B, Sachin G, Padmaja G, and Shivraj S (2019). Importance and Proven Benefits of Ginger on Molecular Basis Study: Review Article. *International Journal of Pharmacy and Biological Sciences*, 9(3):209-219.
- Arnault I, Auger J (2006;). Seleno-compounds in garlic and onion. *J Chromatogr A* 1112: 23-30.
- Ayensu, ES (1978). *Medicinal Plants of West Africa*. Reference Publ. Inc; Algonac, Michigan. 162.
- Backer JA, Klinkenberg D, Wallinga J. (2020) Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20–28 January 2020. *Eur. Surveill*. 25: 2000062.
- Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, Wang M. (2020) Presumed asymptomatic carrier transmission of COVID-19. *JAMA* 323: 1406–1407.
- Beristain-Bauza SDC, Hernández-Carranza P, Cid-Perez TS, Ávila-Sosa R, Ruiz-López II, Ochoa-Velasco CE. (2019). Antimicrobial Activity of Ginger (*Zingiber Officinale*) and Its Application in Food Products. *Food Reviews International*, 35(5): 407-426.
- Bezerra LL, Marinho ES, Marinho MM (2018). Allicin an Inhibiting Potential of HIV Virus: A Molecular Docking Studies Comparative with the Ritonavir® Inhibitor. *International Journal of Recent Research and Review*, 11(3): 6-10
- Bhutta ZA, Black RE, Brown KH, Gardner JM, Gore S, Hidayat A, (1999) Prevention of diarrhea and pneumonia by zinc supplementation in children in developing countries: Pooled analysis of randomized controlled trials. *Zinc Investigators' Collaborative Group. J Pediatr* 135:689–697.

- Bin Sayeed MS, Shams T, Fahim Hossain S, Rahman MR, Mostofa A, Fahim Kadir M. (2014). *Nigella sativa* L. seeds modulate mood, anxiety and cognition in healthy adolescent males. *J Ethnopharmacol.* 152:156–162.
- Bindu A, Anusha H. (2011) Adverse Effects of Highly Active Anti-Retroviral Therapy (HAART). *Journal of Antivirals and Antiretrovirals*, (04):3–3.
- Bourouiba L (2020). Turbulent Gas Clouds and Respiratory Pathogen Emissions: Potential Implications for Reducing Transmission of COVID-19. *JAMA* 323:1837–1838.
- Bradley JM, Organ CL, Lefer DJ. (2016) Garlic-derived organic polysulfides and myocardial protection. *J. Nutr.* 146: 403S–409S.
- Brown KH, Peerson JM, Allen LH. (1998) Effect of zinc supplementation on children's growth: A meta-analysis of intervention trials. In: Sandström B, Walter P, editors. *Role of Trace Elements for Health Promotion and Disease Prevention*. California: S Karger Pub; pp. 76–83.
- Carr AC and Maggini C (2017). Vitamin C and Immune Function. *Nutrients* 9(11): 1211.
- Cava C, Bertoli G, Castiglioni I. (2020) In Silico Discovery of Candidate Drugs against Covid-19. *Viruses*;12: 404.
- Chambial S, Dwivedi S, Shukla KK, John PJ, Sharma P. (2013) Vitamin C in Disease Prevention and Cure: An Overview. *Indian Journal of Clinical Biochemistry*, 28(4): 314-328.
- Chan JFW, Yuan S, Kok KH, To KKW, Chu H, Yang J, Xing F, Liu J, Yip CCY, Poon RWS, Tsoi HW, Lo SKF, Chan KH, Poon VKM, Chan WM, Ip JD, Cai JP, Cheng VCC, Chen H, Hui CKM. (2020) A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: A study of a family cluster. *Lanc* 395: 514–523.
- Charity UO, Uwaifoh A, Charlse II, Williams AA, Kerry EA. (2012) The effect of chronic ingestion of crude *Garcinia kola* on the histology of the liver. *Eur J of Exp Biol* 2: 404-409.
- Chen D, Xu W, Lei Z, Huang Z, Liu J, Gao Z, Peng L. (2020). Recurrence of positive SARS-CoV-2 RNA in COVID-19: A case report. *Int J Infect Dis* 93: 297–299.
- Chen TY, Chen DY, Wen HW, Ou JL, Chiou SS, Chen JM, Wong ML, Hsu WL. (2013). Inhibition of enveloped viruses infectivity by curcumin *PLoS One*. 8:e62482.
- Chinese Centre for Disease Control and Prevention (CCDC). The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19)–China. 2020. Available online: http://weekly.chinacdc.cn/en/article/id/e53946e2-c6c4-41e9-9a9b_fea8db1a8f51 (accessed on 17 February 2020).
- Chun H, Shin DH, Hong BS, Cho WD, Cho HY, Yang HC (2002). Biochemical properties of polysaccharides from black pepper. *Biol Pharm Bull* 25:1203 – 8.
- Chung H.Y. (2019). Role of Garlic and Ginger in Anti-oxidative and Anti-inflammatory Effects in Aging. *Journal of Food Science and Technology*, 4(5):788-795.
- Collise N, Anthony JA, Anna MC, Roland NN. (2011) Crude Ethanolic Extracts of *Garcinia kola* Seeds Heckel (Guttiferae) Prolong the Lag Phase of *Helicobacter pylori*: Inhibitory and Bactericidal Potential. *J of Med Food*. 14: 822-827.
- Correa AD, Jokl L, Carlsson R (1986). Amino acid composition of some *Amaranthus* sp. grain proteins and of its fractions. *Arch Latinoam Nutr* 36:466 – 76.
- Cousins RJ. Zinc. In: Filer LJ, Ziegler EE, editors. (1996) *Present Knowledge in Nutrition*. 7th ed. Washington DC: International Life Science Institute Nutrition Foundation; pp. 293–306.
- Dairaku I, Han Y, Yanaka N, Kato N. (2010). “Inhibitory effect of curcumin on IMP dehydrogenase, the target for anticancer and antiviral chemotherapy agents,” *Bioscience, Biotechnol Biochem*. 74:185–187.
- Dalziel JM. (2012). *The Useful Plants of West Tropical Africa*. Crown Agents for the Colonies, London. 1937. diets containing *Garcinia kola* seed meal. *Rabb Genet* 2: 10-14.
- Diretto G, Rubio-Moraga A, Argandona J, Castillo P, Gomez-Gomez L, Ahrazem O (2017). Tissue-specific accumulation of sulfur compounds and saponins in different parts of garlic cloves from purple and white ecotypes. *Molecules* 22: 1359.
- Doherty VF, Olaniran O, Kanife UC (2010). Antimicrobial activities of *Aframomum melegueta* (alligator pepper). *Int J Biol.* 2: 126-131.
- Durak A, Gawlik-Dziki U, Kowlska I. (2015) coffee with ginger- interactions of biologically active phytochemicals in the model system. *food chemistry*, 166, 261-269.
- Ebana RU, Madunagu BE, Ekpe ED, Otung IN (1991). Microbiological exploitation of cardiac glycosides and alkaloids from *Garcinia kola*, *Borreria ocymoides*, *Kola nitida* and *Citrus aurantifolia*. *J App Bacteriol* 71: 287- 297.
- El-Abhar HS, Abdallah DM, Saleh S. (2003) Gastroprotective effect of black seed. *J.Ethnopharmacol.* 84:251–258.
- Elena SF, Sanjuan R. (2005). Adaptive Value of High Mutation Rates of RNA Viruses: Separating Causes from Consequences. *Journal of Virology*, 79(18):11555–11558.

- El-Halawany AM, Dine RS, El Sayed NS, Hattori M. (2014). Protective effect of *Aframomum melegueta* phenolics against CCl₄-induced rat hepatocytes damage; role of apoptosis and pro-inflammatory cytokines inhibition. *Sci Rep*. 4:5880.
- Elnima EI, Ahmed SA, Mekkawi AG, Mossa JS. (1983). The antimicrobial activity of garlic and onion extracts. *Pharmazie* 38: 747-748.
- Esimone CO, Nwafor SV, Okoli CO, Chah KF, Uzuegbu DB, Chibundu C, Eche MA, Adikwu MU (2002). In Vivo Evaluation of Interaction Between Aqueous Seed Extract of *Garcinia kola* Heckel and Ciprofloxacin Hydrochloride. *Am J of Therap*. 9: 275-280.
- Farombi EO, Owoeye O. (2011). Antioxidant and chemopreventive properties of *Vernonia amygdalina* and *Garcinia biflavonoid*. *Int J of Environ Res Pub Health*. 8:2533–2555.
- Fehr AR, Perlman S. (2015). Coronaviruses: An Overview of Their Replication and Pathogenesis. In: Maier H, Bickerton E, Britton P (eds) *Coronaviruses. Methods in Molecular Biology*, vol 1282. Humana Press, New York, NY.
- Gabriel FI, Emmanuel OO (2011). Pharmacological evaluation of *Garcinia kola* Pharmacological evaluation of *Garcinia kola* nut for antitrichomonal activity. *Int J Pharma Bio Sci* 2: 264-269.
- Gallagher TM, Buchmeier M. (2001) Coronavirus spike proteins in viral entry and pathogenesis. *J. Virol*. 279:371-374.
- Gheznavi, Ghaznavi K. *Tibb-i-Nabvi* (1988) aur jadeed Science. Faisal Publishers, Urdu Baza. Lahore, 1(228): 231-232
- Ghishan FK. (1984) Transport of electrolytes, water, and glucose in zinc deficiency. *J Pediatr Gastroenterol Nutr* 3:608–612.
- Graefe EU, Wittig J, Mueller S, Riethling AK, Uehleke B, Drewelow B, Pforte H, Jacobasch G, Derendorf H, Veit M. (2001) Pharmacokinetics and bioavailability of quercetin glycosides in humans. *J Clin Pharmacol* 41: 492-499.
- Griffiths G, Trueman L, Crowther T, Thomas B, Smith B. (2002) Onions—a global benefit to health. *Phytother Res*. 16: 603-615.
- Guan W, Ni Z, Hu Yu, Liang W, Ou C, He J, Liu L, Shan H, Lei C, Hui, DSC, Du B, Li L, Zeng G, Yuen KY, Chen R, Tang C, Wang T, Chen P, Xiang J, Li S, Wang G, Liu J, Chen J, Li G, Zheng Z, Qiu S, Luo J, Ye C, Zhu S, Zhong N. (2020) Clinical characteristics of 2019 novel coronavirus infection in China. *N. Engl. J. Med*. 382: 1708-1720.
- Hall A, Troupin A, Londono-Renteria B, Colpitts T. (1992) Garlic Organosulfur Compounds Reduce Inflammation and Oxidative Stress during Dengue Virus Infection. *Viruses*, 2017;9(7):159–159.
- Harazem R, Rahman S, Kenawy A. (2019) Evaluation of Antiviral Activity of *Allium Cepa* and *Allium Sativum* Extracts against Newcastle Disease Virus. *Alexandria Journal of Veterinary Sciences*, 61(1): 108-118.
- Haworth WN, Hirst EL. (1933). Synthesis of ascorbic acid. *J Soc Chem Ind (Lond)*, 52:645–647.
- Ho WY, Liang WS, Yeap SK, Beh BK, Younsr AHN. (2012). In vitro and in vivo antioxidant activity of *Vernonia amygdalina* water extracts, *Afric J Biotech*. 11:4090–4094
- Holmes KV (2003). SARS coronavirus: a new challenge for prevention and therapy. *J. Clin. Invest*. 111:1605-1609.
- Hosseinzadeh H, Eskandari M, Ziaee T (2008). Anti-tussive effect of thymoquinone a constituent of *Nigella Sativa* Seeds in guinea pigs. *Pharmacology online*. 2:480–484.
- Hosseinzadeh H, Moghim FF, Mansouri SMT. (2007) Effect of *Nigella sativa* seed extracts on ischemia-reperfusion in rat skeletal muscle. *Pharmacology online*. 2:326–335.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo Li, Xie J, Wang G, Jiang R, Gao Z, Jin Q. (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395: 497–506.
- Huentelman MJ, Zubcevic J, Hernandez Prada JAH Xiao X, Dimitrov DS, Raizada MK, Ostrov DA. (2004) Structure-based discovery of a novel angiotensin-converting enzyme 2 inhibitor. *Hypertension* 44(6): 903–906.
- Igile GO, Oleszek W, Jurzysta M, Burda S, Fanfunso M, Fasanmade AA. (1994) Flavonoids from *Vernonia amygdalina* and their antioxidant activities. *J Agric Food Chem*. 42:2445–2448.
- Ilic NM, Dey M, Poulev AA, Logendra S, Kuhn PE, Raskin I. (2014) Anti-inflammatory activity of grains of paradise (*Aframomum melegueta* Schum) extract. *J of Agr Food Chem*. 62:10452-10457
- Ilkhanizadeh B, Shirpoor A, Ansari MK, Nemati S, Rasmi Y. (2016). Protective Effects of Ginger (*Zingiber officinale*) Extract against Diabetes-Induced Heart Abnormality in Rats. *Diabetes Metab J*.40(1): 46–53.
- Iwu MM, Igboko OA, Okunji CO, Tempesta MS. (1990). Antidiabetic and aldose reductase activities of biflavanones of *Garcinia kola*. *J Pharm Pharmacol*. 42: 290-292.
- Iwu MM. (1989). Food for medicine (Ed.) M. Iwu. In: *Dietary Plants and Masticatories as Sources of Biologically Active Substances*. University of Ife, Nigeria. Ife Press. 303–310.

- Iwu, M. (1993) Handbook of African medicinal plants. CRC Press, Boca Raton, FL., 12: 32-38.
- Jassim SA, Naji MA. (2003). "Novel antiviral agents: a medicinal plant perspective," J App Microbiol. 95:412–427.
- Juliana A, Moctar S, Christopher K, (2006). *Garcinia kola* Heckel. Forest & Landscape Denmark..
- Kakjing DF, Mary OU Nkoli LN. (2014). Anti-pyretic Activity of *Garcinia kola* Seed Extract. Eur J Med Plants 4: 511-521.
- Khaerunnisa S, Kurniawan H, Awaluddin R, Suhartati S, Soetjipto S. (2019) Potential Inhibitor of COVID-19 Main Protease (M^{Pro}) from Several Medicinal Plant Compounds by Molecular Docking Study. Preprints, 1-14
- Khan MR. (1999) Chemical composition and medicinal properties of *Nigella sativa*. Linn. Inflammopharmacology. 7:13–35.
- Khare C.P. (2004) Encyclopedia of Indian medicinal plants. New York: Springer-Verlag Berlin Heidelberg. King JC, Cousins RJ. Zinc. (2006) In: Shils ME, Shike M, Ross AC, Caballero B, Cousins RJ, editors. Modern Nutrition in Health and Disease. 10th ed. Baltimore: Lippincott Williams and Wilkins; pp. 271–285.
- Lai ST. (2005). Treatment of severe acute respiratory syndrome. Eur. J. Clin. Microbiol. Infect. Dis. 24:583-591.
- Lanzotti V (2006). The analysis of onion and garlic. J Chromatogr A 1112: 3-22.
- Lauer AC, Groth N, Haag SF, Darvin ME, Lademann J, Meinke MC. (2013) Dose-dependent vitamin C uptake and radical scavenging activity in human skin measured with in vivo electron paramagnetic resonance spectroscopy. Skin Pharmacol. Physiol. 26:147–154.
- Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, Azman AS, Reich NG, Lessler J. (2020). The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: Estimation and application. Ann. Intern. Med. 172: 577-582.
- Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, Cummings DA. (2009). Incubation periods of acute respiratory viral infections: A systematic review. Lancet Infect. Dis. 9: 291–300.
- Li F. (2014). Receptor Recognition Mechanisms of Coronaviruses: a Decade of Structural Studies. Journal of Virology, 89(4): 1954-1964.
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Leung KS, Lau EH, Wong JY, Xing X, Xiang N, Wu Y, Li C, Chen Q, Li D, Liu T, Zhao J, Liu M, Tu W, Chen C, Jin L, Yang R, Wang Q, Zhou S, Wang R, Liu H, Luo Y, Liu Y, Shao G, Li H, Tao Z, Yang Y, Deng Z, Liu B, Ma Z, Zhang Y, Shi G, Lam TTY, Wu JT, Gao GF, Cowling BJ, Yang B, Leung GM, Feng Z. (2020). Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. N Engl J Med 382: 1199–1207.
- Li S, Tang Z, Li Z, Liu X. (2020). Searching therapeutic strategy of new coronavirus pneumonia from angiotensin-converting enzyme 2: the target of COVID-19 and SARS-CoV. European Journal of Clinical Microbiology & Infectious Diseases, 39:1021–1026.
- Liu S, Xiao G, Chen Y, He Y, Niu J, Escalante CR, Xiong H, Farmer J, Debnath AK, Tien P, Jiang S (2009). Interaction between heptad repeat 1 and 2 regions in spike protein of SARS-associated coronavirus: implications for virus fusogenic mechanism and identification of fusion inhibitors. Lancet 363: 938–947.
- Liu Y, Ning Z, Chen Y, Guo M, Liu Y, Gali NK, Sun L, Duan Y, Cai J, Westerdahl D, Liu X, Xu K, Ho K, Kan H, Fu Q, Lan K. (2020). Aerodynamic Characteristics and RNA Concentration of SARS-CoV-2 Aerosol in Wuhan Hospitals during COVID-19 Outbreak. Nat 582: 557–560
- Lu L, Liu Q, Zhu Y, Chan KH, Qin L, Li Y, Wang Q, Chan JFW, Du L, Yu F, Ma C, Ye S, Yuen KW, Zhang R, Jiang S. (2014). Structure-based discovery of Middle East respiratory syndrome coronavirus fusion inhibitor. Nat. Commun. 5: 3067.
- Lykkesfeldt J, Michels AJ, Frei B. (2014). Vitamin C. Advances in Nutrition 5(1): 16–18.
- Maginnis MS. (2018). Virus–Receptor Interactions: The Key to Cellular Invasion. J Mol Biol. 430(17): 2590–2611.
- Majdalawieh AF, Hmaidan R, Carr RI. (2010). *Nigella sativa* modulates splenocyte proliferation, Th1/Th2 cytokine profile, macrophage function and NK anti-tumor activity. J Ethnopharmacol. 131:268–275.
- Mathew OW, Blessing CD. (2007). Hepatoprotective effects of *Garcinia kola* seed against hepatotoxicity induced by carbon tetrachloride in rats. Nig Soc Exp Biol 4: 17-21.
- Mehrbod P, Aini I, Amini, E, Eslami, M, Torabi, A, Bande, F, Kheiri, M. T. (2013). Assessment of direct immunofluorescence assay in detection of antiviral effect of garlic extract on influenza virus. African Journal of Microbiology Research, 7(21): 2608-2612.
- Mele M.A. (2019). Bioactive compounds and biological activity of ginger. Journal of Multidisciplinary Sciences, 1(1): 1-7.
- Mohammed A, Gbonjubola VA, Koorbanally NA, Islam MS. (2017). Inhibition of key enzymes linked to type 2 diabetes by compounds isolated from *Aframomum melegueta* fruit. Pharm Biol. 55:1010-1016.
- Morabandza CJ, Ongoka RP, Matini L, Epa C, Nkounkou LC, Abena AA. (2013). Chemical Composition of the Mesocarp of *Garcinia kola* Heckel (Clusiaceae) Fruit. Res J of Rec Sci. 2: 53-58.

- Mounce BC, Cesaro T, Carrau L, Vallet T, Vignuzzi M. (2017). Curcumin inhibits Zika and chikungunya virus infection by inhibiting cell binding. *Antiviral Res.* 142:148-157.
- Munaya C (2013). Bitter leaf-based extracts cures hepatitis co-infection and others. *The Guardian Newspaper*, July, 25, 2013.
- Munster VJ, Koopmans M, van Doremalen N, van Riel D, de Wit E. (2020). A novel coronavirus emerging in China—key questions for impact assessment. *N Engl J Med* 2020; 382: 692–694. *Pathogens* 9: 692–694.
- Nathan M, Scholten R. (1999). The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines. *Ann Int Med* 130: 459.
- Navarro JA, del Pozo MD, Gastaminza G, Moneo I, Audicana MT, Fernández de Corres L. (1995). *Allium cepa* seeds: a new occupational allergen. *J Allergy Clin Immunol.* 96: 690-693.
- Nemeth K, Piskula MK. (2007). Food content, processing, absorption and metabolism of onion flavonoids. *Crit Rev Food Sci Nutr* 47: 397-409.
- Nwauzoma AB, Dappa MS. (2013) Ethnobotanical Studies of Port Harcourt Metropolis, Nigeria. *International Scholarly Research Notices*: 1-11.
- Nwokochaa CR, Owua DU, Ufearob CS Iwualac MOE. (2011). Comparative study on the efficacy of *Garcinia kola* in reducing some heavy metal accumulation in liver of Wistar rats. *J of Ethnopharmacol.* 135: 488–491.
- Ofokansi KC, Mbanefo AN, Ofokansi MN, Esimone CO. (2008). Antibacterial Interaction of Crude Methanol Extract of *Garcinia kola* Seed with Gatifloxacin. *Trop J of Pharm Res* 7: 1159- 1165.
- Okoko T, Ndoni S. (2010). The effect of *Garcinia kola* extract on lipopolysaccharide induced tissue damage in rats. *Trop J Pharm Res* 17: 88–93.
- Okoli CO, Akah PA, Nwafor SV, Ihemelandu UU, Amadife C. (2007). Anti-inflammatory activity of seed extracts of *Aframomum melegueta*. *J of Herbs, Spices Med Plants.* 13:11-21
- Okwu DE (2005). Phytochemicals, vitamins and mineral contents of two Nigerian medicinal plants. *Int J Mol Med Adv Sci.* 1:375-381.
- Omar A, Ghosheh S, Abdulghani A, Houdi A, Crookscor PA. (1999). High performance liquid chromatographic analysis of the pharmacologically active quinones and related compounds in the oil of the black seed (*Nigella sativa* L). *J Pharm Biomed Anal* 19:757 – 62
- Omosa LK, Midiwo JO, Kuete V. (2017). “Curcuma longa,” in *Therapeutic Potential Against Metabolic, Inflammatory, Infectious and Systemic Diseases*, ed. (Cambridge, MA: Academic Press) :425–435.
- Onasanwo SA, Singh N, Olaleye SB, Palit G. (1990). Anti-ulcerogenic and proton paracetamol-induced hepatotoxicity in rats. *J of ethnopharmacol* 29: 207–211.
- Osuntokun OT, Oluduro AO, Idowu TO, Omotuyi AO. (2017). Assessment of Nephro toxicity, Anti-inflammatory and Antioxidant properties of Epigallocatechin, Epicatechin and Stigmasterol phytosterol (synergy) Derived from ethyl acetate stem bark extract of *Spondias mombin* on Wistar Rats Using Molecular method of analysis. *J of Mol Microbiol.* 1:1-11.
- Otor JU, Abdulkadir U, Abu M.A. (2001). Some biological activities of *Garcinia kola* in growing rats. *Vet arhiv* 71: 287- 297.
- Penduka D, Okoh OO, Okoh AI. (2011). In Vitro Antagonistic Characteristics of Crude Aqueous and Methanolic Extracts of *Garcinia kola* (Heckel) Seeds against Some *Vibrio* Bacteria. *Mol.* 16: 2754-2765.
- Prabakaran P, Xiao X, Dimitrov DS. (2004). A model of the ACE2 structure and function as a SARS-CoV receptor. *Biochem Biophys Res Commun* 314: 235–241.
- Prasad AS, Miale A, Jr, Farid Z, Sandstead HH, Schulert AR. (1963). Zinc metabolism in patients with the syndrome of iron deficiency anemia, hepatosplenomegaly, dwarfism, and hypogonadism. *J Lab Clin Med.* 61:537–549
- Raj VS, Mou H, Smits SL, Dekkers DHW, Müller MA, Dijkman R, Muth D, Demmers JAA, Zaki A, Fouchier RAM, Thiel V, Drosten C, Rottier PJM, Osterhaus ADME, Bosch BJ and Haagmans BL. (2013) Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus. *EMC. Nature* 495:251–254.
- Reguera J, Santiago C, Mudgal G, Ordoño D, Enjuanes L, and Casasnovas JM. (2012). Structural Bases of Coronavirus Attachment to Host Aminopeptidase N and Its Inhibition by Neutralizing Antibodies. *PLoS Pathog.* 8(8): e1002859.
- Rose P, Whiteman M, Moore PK, Zhu YZ. (2005). Bioactive S-alk(en)yl cysteine sulfoxide metabolites in the genus *Allium*: the chemistry of potential therapeutic agents. *Nat Prod Rep* 22: 351-368.
- Sahardi NFN and Makpol S. (2019). Ginger (*Zingiber officinale* Roscoe) in the Prevention of Ageing and Degenerative Diseases: Review of Current Evidence. *Evidence-Based Complementary and Alternative Medicine*, Vol. 5054395: 1-7.

- Sahin K, Kilic E, Balcikanli Z, Ates N, Orhan C, Tuzcu M, Juturu V. (2019). Ginger provides neuroprotection in experimental model of traumatic brain injury. *Biochemistry and Molecular Biology*, 33(1): Pages 1b1-872.4.
- Salem ML. (2005). Immunomodulatory and therapeutic properties of the *Nigella sativa* L. seed. *Int Immunopharmacol*. 5:1749–1770.
- Sandrine Belouzard-, Jean K Millet, Beth N Licitra, Gary R Whittaker. (2012). Mechanisms of Coronavirus Cell Entry Mediated by the Viral Spike Protein. *Viruses*, 4(6):1011-1033.
- Sandstead HH, Prasad AS, Schulert AR, Farid Z, Miale A Jr, Bassilly S, Darby WJ. (1967). Human zinc deficiency, endocrine manifestations and response to treatment. *Am J Clin Nutr*. 20:422–442.
- Schlueter AK and Johnston CS. (2010). Vitamin C: Overview and Update. *Journal of Evidence-Based Complementary and Alternative Medicine*, 16(1): 49-57.
- Seanego CT, Ndip RN. (2012). Identification and Antibacterial Evaluation of Bioactive Compounds from *Garcinia kola* (Heckel) Seeds *Mol*. 17: 6569-6584.
- Shafi G, Munshi A, Hasan TN, Alshatwi AA, Jyothy A, David KY. (2009). Anti cancer activity of *Nigella sativa*. *Cancer Cell Int*. 9:29.
- Shang A, Cao SY, Xu XY, Gan RY, Mavumengwana V, Corke H. (2019). Bioactive Compounds and Biological Functions of Garlic (*Allium sativum* L.). *Foods*, 8: 246
- Shankar AH, Genton B, Baisor M, Paino J, Tamja S, Adiguma T, Wu L, Rare L, Bannon D, Tielsch JM, West KP Jr, Alpers MP. (2000). The influence of zinc supplementation on morbidity due to *Plasmodium falciparum*: A randomized trial in preschool children in Papua New Guinea. *Am J Trop Med Hyg*. 62:663–669.
- Sharma N. (2019). Efficacy of Garlic and Onion against virus. *International Journal in Research and Pharmaceutical Sciences*, 4(10): 3578-3586.
- Sharma P, Raghavan SA, Saini R, Dikshit M. (2004). Ascorbate-mediated enhancement of reactive oxygen species generation from polymorphonuclear leukocytes: Modulatory effect of nitric oxide. *J. Leukoc. Biol*. 75:1070–1078.
- Shojai TM, Langeroudi AG, Karimi V, Barin A, Sadri N. (2016). The effect of *Allium sativum* (Garlic) extract on infectious bronchitis virus in specific pathogen free embryonic egg. *Avicenna Journal of Phytomedicine*, 6(4):267–458.
- Singh R and Singh K (2019). *Zingiber Officinale*: A Spice with multiple roles. *Research Journal of Life Sciences, Bioinformatics, Pharmaceutical and Chemical Sciences*, 5(2):113.
- Spellberg B, Haddix M, Lee R, Butler-Wu S, Holtom P, Yee H, Gounder P. (2020). Community Prevalence of SARS-CoV-2 Among Patients With Influenza like Illnesses Presenting to a Los Angeles Medical Center in March 2020. *JAMA* 323: 1966-1967.
- Stadler K, Masignani V, Eickmann M, Becker S, Abrignani S, Klenk HD, Rappuoli R (2003). SARS-beginning to understand a new virus. *Nat. Rev. Microbiol*. 1:209-218.
- Suleria HA, Butt MS, Anjum FM, Saeed F, Khalid N. (2015). Onion: nature protection against physiological threats. *Crit Rev Food Sci Nutr* 55: 50-66.
- Szychowski KA, Rybczynska-Tkaczyk K, Gawel-Beben K, Swieca M, Karas M, Jakubczyk A, Matysiak M, Binduga UE, Gminski J. (2018). Characterization of active compounds of different garlic (*Allium sativum* L.) cultivars. *Pol. J. Food Nutr. Sci*. 68: 73–81.
- Tanaka K, Arita M, Sakurai H, Ono N, Tezuka Y. (2015). Analysis of Chemical Properties of Edible and Medicinal Ginger by Metabolomics Approach. *BioMed Research International*, vol Article ID 671058: 1-7.
- Tebekeme, O. (2009). In vitro antioxidant and free radical scavenging activities of *Garcinia kola* seeds. *Food and Chem Toxicol*. 47: 2620–2623.
- Teissier E, Penin F, Pécheur EI. (2011). Targeting Cell Entry of Enveloped Viruses as an Antiviral Strategy. *Molecules*, 16(1): 221–250.
- Terashima K, Takaya Y, Niwa M. (2002). Powerful antioxidative agents based on garcinoic acid from *Garcinia kola*. *Bioorganic Med Chem* 10: 1619-1625.
- Tubek S. (2007). Selected zinc metabolism parameters in premenopausal and postmenopausal women with moderate and severe primary arterial hypertension. *Biol Trace Elem Res*. 116:249–256.
- Turnlund JR, King JC, Keyes WR, Gong B, Michel MC. (1984). A stable isotope study of zinc absorption in young men: Effects of phytate and alpha-cellulose. *Am J Clin Nutr* 40:1071-1077.
- Udochukwu U, Omeje FI, Uloma IS, Oseiwe ID. (2015). Phytochemical Analysis of *Vernonia Amygdalina* and *Ocimum Gratissimum* Extracts and their antibacterial activity on some Drug Resistant Bacteria. *Am J of Res Comm*. 3:225-235.
- Unigwe CR, Nwakpu PE. (2009). Effect of ingestion of *Garcinia kola* seed on Erythrocytes of Wistar Rats. *Nig J Physiol Sci* 20: 30-32.

- Valacchi G, Muresan XM, Sticozzi C, Belmonte G, Pecorelli A, Cervellati F, Demaude J, Krol Y, Oresajo C. (2016). Ozone-induced damage in 3D-skin model is prevented by topical vitamin C and vitamin E compound mixtures application. *J. Dermatol. Sci.* 82:209–212.
- Valdés F. (2006). Vitamin C. *Actas Dermosifiliogr*, 97(9): 557-568.
- Veenemans J, Milligan P, Prentice AM, Schouten LR, Inja N, van der Heijden AC, de Boer LC, Jansen EJ, Koopmans AE, Enthoven WT, Kraaijenhagen RJ, Demir AY, Uges DR, Mbugi EV, Savelkoul HF, Verhoef H. (2011). Effect of supplementation with zinc and other micronutrients on malaria in Tanzanian children: A randomised trial. *PLoS Med.* 8:e1001125.
- Venugopal R, Liu RH. (2012). Phytochemicals in diets for breast cancer prevention: The importance of resveratrol and ursolic acid. *F Sci Hum Well.* 1: 1-13.
- Wang L, Junker D, Collisson EW. (1993). Evidence of natural recombination within the S1 gene of infectious bronchitis virus. *Virology* 192:710–716.
- Wang YC, Guan M, Zhao X, Li XL. (2018). Effects of garlic polysaccharide on alcoholic liver fibrosis and intestinal microflora in mice. *Pharm. Biol.* 56: 325–332.
- Weber ND, Andersen DO, North JA, Murray BK, Lawson LD, Hughes BG. (1992). In vitro virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta Med.* 58(5):417-423.
- WHO, (1999) on *Allium cepa* dosing Press release.
- WHO. Director-General's Opening Remarks at the Media Briefing on COVID-19–3 March 2020; World Health Organization: Geneva, Switzerland, 15 July 2020.
- WHO. Statement on the Second Meeting of the International Health Regulations Emergency Committee Regarding the Outbreak of Novel Coronavirus (2019-nCoV); World Health Organization: Geneva, Switzerland, 30 January 2020. Available online: [https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)) (accessed on 3 March 2020).
- Wintergerst ES, Maggini S, Hornig DH. (2006). Immune-enhancing Role of Vitamin C and Zinc and Effect on Clinical Conditions. *Ann Nutr Metab*, 50(2):85-94.
- Wong CK, Lam CWK, Wu AKL, Ip WK, Lee NLS, Chan IHS, Lit LCW, Hui DSC, Chan MHM, Chung SSC, Sung JJY. (2004). Plasma inflammatory cytokines and chemokines in severe acute respiratory syndrome. *Clin Exp Immunol* 136: 95–103.
- Xia J, Tong J, Liu M, Shen Y, Guo D. (2020). Evaluation of coronavirus in tears and conjunctival secretions of patients with SARS-CoV-2 infection. *J Med Viro* 92 :589-594.
- Xia Shuai, Lei Yan, Wei Xu, Anurodh Shankar Agrawal, Abdullah Algaissi, Chien-Te K. Tseng, Qian Wang, Lanying Du, Wenjie Tan, Ian A. Wilson, Shibo Jiang, Bei Yang, Lu Lu. (2019). A pan-coronavirus fusion inhibitor targeting the HR1 domain of human coronavirus spike. *Science Advances*, 5: 1-15.
- Yang C (2020). Does hand hygiene reduce SARS-CoV-2 transmission? *Graefes Arch Clin Exp Ophthalmol* 258: 1133-1134.
- Zhang JJ, Xin Shen, Yan YM, Wang Y, Cheng YX (2019). Discovery of anti-SARS-CoV-2 agents from commercially available flavor via docking screening. 12;132-38
- Zhao Y, Zhao Z, Wang Y, Zhou Y, Ma Y, Zuo W. (2020). Single-cell RNA expression profiling of ACE2, the putative receptor of Wuhan 2019-nCoV. *BioRxiv* 1-13.
- Zohri AN, Abdel-Gawad K, Saber S. (1995). Antibacterial, antidermatophytic and antitoxigenic activities of onion (*Allium cepa* L.) oil. *Microbiol Res* 150: 167-172.